

Stability and collapse of a hybrid Bose-Einstein condensate of atoms and molecules

Sadhan K Adhikari

Instituto de Física Teórica, Universidade Estadual Paulista, 01.405-900 São Paulo, São Paulo, Brazil

(February 1, 2008)

Abstract

The dynamics of stability and collapse of a trapped atomic Bose-Einstein condensate (BEC) coupled to a molecular one is studied using the time-dependent Gross-Pitaevskii (GP) equation including a nonlinear interaction term which can transform two atoms into a molecule and vice versa. We find interesting oscillation of the number of atoms and molecules for a BEC of fixed mass. This oscillation is a consequence of continuous transformation in the condensate of two atoms into a molecule and vice versa. For the study of collapse an absorptive contact interaction and an imaginary quartic three-body recombination term are included in the GP equation. It is possible to have a collapse of one or both components when one or more of the nonlinear terms in the GP equation are attractive in nature, respectively.

PACS Number(s): 03.75.Fi, 05.30.Jp

I. INTRODUCTION

The experimental detection [1–3] of Bose-Einstein condensation (BEC) at ultralow temperature in dilute trapped alkali metal and hydrogen atoms has initiated intense theoretical activities in this subject [4–14]. Many properties of the condensate are usually described by the nonlinear meanfield Gross-Pitaevskii (GP) equation [15] which can be considered to be a generalization of the Schrödinger equation in the presence of interatomic interaction leading to a nonlinear term.

Two interesting features of BEC are (a) the collapse in the case of an attractive atomic interaction [3,4] and (b) the possibility of the formation of a hybrid atomic and molecular condensate [6–10] coupled by coherent Raman transition [16] or Feshbach resonance [8,17]. Although there has been experimental observation of a coupled BEC involving two atomic states [18], so far there has been no experimental realization of a molecular BEC. Because of the intrinsic interest in a hybrid atomic and molecular BEC, in this work the dynamics of such a BEC is studied using the time-dependent GP equation with special attention to its stability and collapse. For the case of a trapped two-species BEC a large repulsive interaction between the two species can induce instability in the system [19].

For attractive interatomic interaction [3,4], the condensate is stable for a maximum critical number of atoms. When the number of atoms increases beyond the critical number, due to interatomic attraction the radius of BEC tends to zero and the central density tends to infinity. Consequently, the condensate collapses emitting atoms until the number of atoms is reduced below the critical number and a stable configuration is reached. With a supply of atoms from an external source the condensate can grow again and thus a series of collapses can take place, which was observed experimentally in the BEC of ^7Li with attractive interaction [3,4]. Theoretical analyses based on the GP equation also confirm the collapse [3,4,14].

The possibility of the formation of a hybrid trapped atomic BEC coupled to a BEC of diatomic molecules via either Raman photoassociation [16] or a Feshbach resonance [8,17]

has been suggested and studied recently [6,7]. These authors used a nonlinear interaction term [6,20] in the Hamiltonian that allows for a coherent number-conserving nondissipative transition of two atoms to a molecule and vice versa. Possible candidates for molecular BEC are $^{87}\text{Rb}_2$, $^{85}\text{Rb}_2$, $^{23}\text{Na}_2$ etc [7,21]. The coherently coupled BEC could allow localized solitons even in the absence of a trap [6]. However, in this work we shall only consider the interesting dynamics of this hybrid BEC in the presence of a trap. Because of the general interest there have been several theoretical studies involving two coupled atomic condensates [22]

Here we critically study the atomic plus molecular hybrid BEC in the presence of attractive bosonic interaction(s). The usual GP equation allowing for transition of two atoms to a molecule and vice versa conserves the overall number of atoms. In the study of the dynamics of this BEC, oscillation of the individual numbers of atoms and molecules are found with the conservation of overall total number (number of atoms plus twice the number of diatomic molecules). The dynamics of collapse for attractive interaction is studied by introducing absorptive contact interactions which allow for a growth in the particle numbers from an external source in addition to imaginary quartic three-body interactions responsible for recombination loss [4]. In the presence of the imaginary contact and three-body interactions the GP equation does not conserve the overall number of atoms. If the strengths (and signs) of these two imaginary interactions are properly chosen, the solution of the GP equation could produce a growth of the condensate with time for numbers of atoms and molecules less than the respective critical numbers. Once they increase past these critical numbers the three-body interaction takes control and these numbers suddenly drop below the critical level by recombination loss signaling a collapse [4]. Then the absorptive term takes over and the individual numbers start to increase. This continues indefinitely showing an infinite sequence of collapses.

There are three types of nonlinear interactions in the GP equation for the hybrid BEC of atoms and molecules: e.g., between two atoms, between two molecules, and between an atom and a molecule. If one of these is attractive, one component of the condensate

may experience collapse. If two are attractive one can have collapse in both components. Specifically, one can also have collapse of both components if only the interaction between an atom and a molecule is sufficiently attractive.

In Sec. II we present a set of coupled GP equations which is appropriate for the study of the hybrid atomic-molecular BEC including a new nonlinear interaction term which allows for the possibility of transformation of two atoms in the condensate to a molecule and vice versa. In addition we include an absorptive contact term and a three-body recombination term appropriate for studying collapse of the BEC. In Sec. III we solve the coupled GP equation numerically and study the hybrid atomic-molecular BEC under different possibilities of atomic and molecular interactions. Finally, in Sec. IV we give some concluding remarks.

II. COUPLED GROSS-PITAEVSKII EQUATION FOR ATOMIC-MOLECULAR CONDENSATE

Here, in addition to the kinetic energies, trap potentials, and usual nonlinear interaction terms the following interaction responsible for a nondissipative transfer of two atoms into a molecule and vice versa is also considered in the hamiltonian [6,20]

$$H_{\text{int}} = \frac{\chi}{2} \int d\mathbf{r} \left[\hat{\psi}_1^2 \hat{\psi}_2^\dagger + \hat{\psi}_1^\dagger{}^2 \hat{\psi}_2 \right], \quad (1)$$

where $\hat{\psi}_1$ ($\hat{\psi}_2$) is the complex atomic (diatomic molecular) field and χ the coupling. For a nonzero χ neither the number of atoms nor that of the molecules is conserved. However the total mass of all the atoms and molecules is conserved. In a BEC the chemical conversion of two atoms to a diatomic molecule is dominated by coherent stimulated emission, in which transitions are enhanced by the number of molecules already occupying the ground state. The conversion of atoms to molecules could be resonant via a Feshbach resonance. Or, it could be effected via stimulated Raman coupling introduced by two laser fields. In this fashion it is possible to select a specific molecular state in the condensate.

In the presence of the extra interaction (1) the mean-field GP equation for wave-function components become [6,7]

$$\left[-\frac{\hbar^2}{2m} \frac{1}{r} \frac{\partial^2}{\partial r^2} r + \frac{1}{2} m \omega^2 r^2 + \sum_{j=1}^2 g_{1j} |\psi_j(r, \tau)|^2 - i\hbar \frac{\partial}{\partial \tau} \right] \psi_1(r, \tau) + \chi \psi_2(r, \tau) \psi_1^*(r, \tau) = 0, \quad (2)$$

$$\left[-\frac{\hbar^2}{4m} \frac{1}{r} \frac{\partial^2}{\partial r^2} r + c m \omega^2 r^2 + \sum_{j=1}^2 g_{2j} |\psi_j(r, \tau)|^2 - i\hbar \frac{\partial}{\partial \tau} \right] \psi_2(r, \tau) + \frac{\chi}{2} \psi_1^2(r, \tau) = 0, \quad (3)$$

where m is the atomic mass, $2m$ the diatomic molecular mass, g_{ij} the nonlinear interaction, ω the frequency of the harmonic oscillator trap, and $i = 1$ (2) denotes an atom (molecule). The parameter c has been introduced in (3) to modify the frequency of the trap for molecules. The diagonal interaction term g_{ii} is related to the scattering length a_i of i -type bosons via $g_{ii} = 4\pi\hbar^2 a_i / m_i$, where m_i is the corresponding mass. The wave-function components are normalized as

$$\int d^3r |\psi_1(r, \tau)|^2 = N_1, \quad (4)$$

$$\int d^3r |\psi_2(r, \tau)|^2 = N_2, \quad (5)$$

where N_j is the number of j -type condensed boson, For $\chi = 0$ the number of bosons of each type is conserved. For $\chi \neq 0$, the number of bosons of each type is not separately conserved and there will be continuous transfer of one type of bosons to another and vice versa. However, the overall total number ($N_1 + 2N_2$) is conserved.

As in Refs. [13] it is convenient to use dimensionless variables defined by $x = \sqrt{2}r/a_{\text{ho}}$, $t = \tau\omega$, and $\phi_i(x, t) = x\psi_i(r, \tau)(\sqrt{2\pi}a_{\text{ho}}^3)^{1/2}$ where $a_{\text{ho}} \equiv \sqrt{\hbar/(m\omega)}$. In terms of these, (2) and (3) become

$$\left[-\frac{\partial^2}{\partial x^2} + \frac{x^2}{4} + \sum_{j=1}^2 n_{1j} \frac{|\phi_j(x, t)|^2}{x^2} - i\xi_1 \frac{|\phi_1(x, t)|^4}{x^4} + i\gamma_1 - i\frac{\partial}{\partial t} \right] \phi_1(x, t) + \eta \frac{\phi_2(x, t)\phi_1^*(x, t)}{x} = 0, \quad (6)$$

$$\left[-\frac{1}{2}\frac{\partial^2}{\partial x^2} + \frac{cx^2}{2} + \sum_{j=1}^2 n_{2j} \frac{|\phi_j(x,t)|^2}{x^2} - i\xi_2 \frac{|\phi_2(x,t)|^4}{x^4} + i\gamma_2 - i\frac{\partial}{\partial t} \right] \phi_2(x,t) + \frac{\eta}{2} \frac{\phi_1^2(x,t)}{x} = 0, \quad (7)$$

where $n_{ij} \equiv g_{ij}m/(\sqrt{2}\pi\hbar^2 a_{\text{ho}})$ is negative (positive) when the corresponding interaction is attractive (repulsive) and the dimensionless coupling $\eta = \chi/[\hbar\omega(\sqrt{2}\pi a_{\text{ho}}^3)^{1/2}]$. The diagonal elements of n_{ij} are $n_{ii} = 2\sqrt{2}(a_i/a_{\text{ho}})$. In (6) and (7) a diagonal absorptive (strength γ_i) and quartic three-body (strength ξ_i) interactions, appropriate to study collapse, are introduced [4]. A nonzero γ_i allows the possibility of the absorption of atoms and molecules from the surrounding into the condensate, whereas a nonzero ξ_i allows for the possibility of ejection of particles from the condensate due to three-body recombination. Consequently, the total mass of the condensate is not conserved for nonzero γ_i and ξ_i . A nonzero γ_i and ξ_i is appropriate for the study of collapse of the system.

The fluctuation of atomic and molecular numbers is best studied via the quantities

$$N_{\text{at}} = \frac{1}{N_1^{(0)}} \int_0^\infty |\phi_1(x,t)|^2 dx, \quad (8)$$

$$N_{\text{mol}} = \frac{1}{N_2^{(0)}} \int_0^\infty |\phi_2(x,t)|^2 dx, \quad (9)$$

where $N_1^{(0)}$ and $N_2^{(0)}$ are the initial numbers of the two types. In the presence of a general absorptive interaction $\eta \neq 0$, $\gamma_i \neq 0$ and $\xi_i \neq 0$, the quantities N_{at} and N_{mol} carry the information about time-evolution of the number of atoms and molecules in the condensate. For $\gamma_i = \xi_i = \eta = 0$, the number of atoms and molecules are separately conserved and one has $N_{\text{at}} = N_{\text{mol}} = 1$. For $\gamma_i = \xi_i = 0$ and $\eta \neq 0$, there is no separate number conservation for atoms and molecules: $N_{\text{at}} \neq 1$, $N_{\text{mol}} \neq 1$; although the total mass of the condensate is conserved. For $\gamma_i \neq 0$ and/or $\xi_i \neq 0$ there is no conservation of the total mass of the condensate.

The root-mean-square (rms) radius $x_{\text{rms}}(t)$ of component i is given by

$$x_{\text{rms}}(t) = \left[\frac{\int_0^\infty x^2 |\phi_i(x,t)|^2 dx}{\int_0^\infty |\phi_i(x,t)|^2 dx} \right]^{1/2}. \quad (10)$$

The oscillation of this rms radius denote radial oscillation of the condensate.

III. NUMERICAL METHOD

To solve (6) and (7) numerically one needs the proper boundary conditions as $x \rightarrow 0$ and ∞ . For a confined condensate, for a sufficiently large x , $\phi_i(x, t)$ must vanish asymptotically. Hence all the nonlinear terms proportional to the product of wave-function components can eventually be neglected in the GP equation for large x . Consequently the asymptotic form of the physically acceptable solution is given by $\lim_{x \rightarrow \infty} |\phi_i(x, t)| \sim \exp(-x^2/4)$. Next we consider (6) and (7) as $x \rightarrow 0$. The nonlinear terms approach a constant in this limit because of the regularity of the wave function at $x = 0$ and one has the condition $|\phi_i(0, t)| = 0$.

Next we discretize (6) and (7) in both space and time by using a space step h and time step Δ with a finite difference scheme using the unknown $(\phi_i)_p^k$ which will be approximation of the exact solution $\phi_i(x_p, t_k)$ where $x_p = ph$ and $t_k = k\Delta$. After discretization we obtain a set of algebraic equations which could then be solved by using the known asymptotic boundary conditions. The numerical procedure is similar to that in refs. [4,13]. The time derivative in these equations involves the wave functions at times t_k and t_{k+1} . As in the uncoupled case we express the wave functions and their derivatives in (6) and (7) in terms of the average over times t_k and t_{k+1} [23] and the resultant scheme leads to accurate results and good convergence. In practice we use the following Crank-Nicholson-type scheme [23] to discretize the partial differential equations (6) and (7)

$$\begin{aligned}
i \frac{(\phi_1)_p^{k+1} - (\phi_1)_p^k}{\Delta} = & -\frac{1}{2h^2} \left[(\phi_1)_{p+1}^{k+1} - 2(\phi_1)_p^{k+1} \right. \\
& \left. + (\phi_1)_{p-1}^{k+1} + (\phi_1)_{p+1}^k - 2(\phi_1)_p^k + (\phi_1)_{p-1}^k \right] \\
& + \frac{1}{2} \left[\frac{x_p^2}{4} + \sum_{l=1}^2 n_{1l} \frac{|(\phi_l)_p^k|^2}{x_p^2} - i\xi_1 \frac{|(\phi_1)_p^k|^4}{x_p^4} + i\gamma_1 \right] \\
& \times \left[(\phi_1)_p^{k+1} + (\phi_1)_p^k \right] + \frac{\eta(\phi_2)_p^k}{x_p} (\phi_1)_p^{k*}, \tag{11}
\end{aligned}$$

$$i \frac{(\phi_2)_p^{k+1} - (\phi_2)_p^k}{\Delta} = -\frac{1}{4h^2} \left[(\phi_2)_{p+1}^{k+1} - 2(\phi_2)_p^{k+1} \right.$$

$$\begin{aligned}
& + (\phi_2)_{p-1}^{k+1} + (\phi_2)_{p+1}^k - 2(\phi_2)_p^k + (\phi_2)_{p-1}^k \Big] \\
& + \frac{1}{2} \left[\frac{cx_p^2}{2} + \sum_{l=1}^2 n_{2l} \frac{|(\phi_l)_p^k|^2}{x_p^2} - i\xi_2 \frac{|(\phi_2)_p^k|^4}{x_p^4} + i\gamma_2 \right] \\
& \times \left[(\phi_2)_p^{k+1} + (\phi_2)_p^k \right] + \frac{\eta(\phi_1)_p^k}{2x_p} (\phi_1)_p^k.
\end{aligned} \tag{12}$$

Considering that the wave function components ϕ_i are known at time t_k , (11) and (12) are equations in the unknowns $(\phi_i)_{p+1}^{k+1}$, $(\phi_i)_p^{k+1}$ and $(\phi_i)_{p-1}^{k+1}$. In a lattice of N points (11) and (12) represent a tridiagonal set for $p = 2, 3, \dots, (N-1)$ for a specific component ϕ_i . This set has a unique solution if the wave functions at the two end points $(\phi_i)_1^{k+1}$ and $(\phi_i)_N^{k+1}$ are known [23]. In the present problem these values at the end points are provided by the known asymptotic boundary conditions. This tridiagonal set of equations is solved by Gaussian elimination and back-substitution [23].

To solve (11) and (12) we employ space step $h = 0.0001$ with $x_{\max} \leq 15$ and time step $\Delta = 0.05$ with $t_{\max} = 100$. After some experimentation we find that these values of the steps give good convergence. The iteration is started with the known normalized (harmonic oscillator) solution of (6) and (7) obtained with $n_{ij} = \gamma_i = \xi_i = \eta = 0$ with normalization

$$\int_0^\infty |\phi_i(x, t)|^2 = N_i^{(0)}, \quad i = 1, 2, \tag{13}$$

where $N_i^{(0)}$ is the initial number of atoms. The nonlinear parameters n_{ij} and η are increased by equal amounts over 500 to 1000 time iterations starting from zero until the desired final values are reached. During this process we keep $\xi_i = \gamma_i = 0$. The resulting solution is the ground state of the condensate corresponding to the specific nonlinearity.

The time-dependent approach is the most suitable for solving time-evolution problems. In the present study we only consider evolution problems starting from a stable condensate at $t = 0$. In these cases the stationary problem is solved first and the wave function so obtained serves as the starting wave function for the time-evolution problem.

To study the effect of an external source of bosons on preformed condensate(s) we need to set $\gamma_i \neq 0$ and $\xi \neq 0$ and perform the time iteration of the GP equations (6) and (7) starting at $t = 0$ from the solution obtained with $\gamma_i = 0$ and $\xi_i = 0$. A nonzero γ_i will

make the condensate grow in size absorbing bosons from outside. In the case of attractive interaction(s), as the condensate grows past the critical size a nonzero ξ will allow for three-body recombination so that the condensate can shrink and experience the so called collapse by emitting particles. Hence a nonzero γ_i and ξ_i could simulate the dynamical growth of the condensate in general as well as its collapse for attractive interaction.

IV. NUMERICAL RESULT

Instead of concentrating on a specific atomic-molecular BEC, here we study some universal features of such a BEC related to stability and collapse. For a specific application we require the knowledge of the different scattering lengths: those for two atoms, two molecules and between an atom and a molecule. At present these scattering lengths are not all known for a specific system.

A. Repulsive Interaction: Stable Condensate

First we consider the nondissipative solutions of (6) and (7) obtained with $\eta \neq 0$ and $\gamma_i = \xi_i = 0$ for all repulsive interactions and demonstrate the oscillation of numbers of atoms and molecules. In this case the condensate increases in size indefinitely with time if a nonzero γ_i and ξ_i are used. This is why it can be termed stable. However, there is spontaneous transformation of atoms to molecules and vice versa even for $\gamma_i = \xi_i = 0$ and the number of atoms and molecules oscillate with time. The numerical results are convergent for small values of nonlinear parameters n_{ij} and η . Here to demonstrate the viability of the present numerical scheme we present results for reasonably large values of n_{ij} . Specifically, we consider $c = 1/16$, $N_1^{(0)} = N_2^{(0)} = 10000$, $n_{11} = n_{22} = 0.001$, $n_{12} = n_{21} = 0.0005$, $x_{\max} = 15$ and $\eta = 0.001, 0.005$, and 0.01 . For a typical experimental situation [11] this corresponds to $2\sqrt{2}|a_i|/a_{\text{ho}} = 10^{-3}$. The solution of the complete coupled GP equations (6) and (7) is obtained by iteration starting with the solution of the uncoupled Schrödinger equation for the harmonic oscillator problem obtained by setting $n_{ij} = \eta = 0$ in these equations.

The nonlinear parameters n_{ij} are increased in small steps during iteration until their final values are reached after 1000 iterations. We then obtain the desired solution of the coupled GP equation. In figure 1 (a) we plot the atomic and molecular wave functions ϕ_1 and ϕ_2 obtained from this procedure.

We continue the time iteration after the solutions of the coupled GP equation shown in figure 1 (a) are obtained at a specific time, which we now call $t = 0$, without changing the nonlinear parameters anymore. In figure 1 (b) we plot the quantities N_{at} and N_{mol} vs. time t for $t > 0$. For $\eta \neq 0$, as shown in figure 1 (b), the numbers N_{at} and N_{mol} undergo rapid oscillations. In figure 1 (c) we plot the corresponding rms radii of the atomic and molecular condensates for $t > 0$. We find that the rms radii also execute oscillation along with N_{at} and N_{mol} due to continued dynamical conversion of atoms to molecules and vice versa. The oscillation of the rms radii is related to the radial oscillation of the condensate.

B. Attractive Interaction: Route to Collapse

Now we consider the case of attractive atomic and molecular interaction(s) which may lead to collapse. We note that η is a nonlinear parameter in the coupled GP equation and a large value of the nonlinear parameter η will imply a large numerical error in the solution. A very small η on the other hand will not give adequate feedback between atoms and molecules. We study collapse in the hybrid BEC under several conditions with $\eta = 0.00625$, as this value of η gives adequate feedback between atoms and molecules and accurate numerical result.

First we study the simplest case of collapse by taking all interactions to be attractive corresponding to a negative n_{ij} for all i and j . Now both components of BEC could experience collapse. In this case the wave function components calculated numerically as in figure 1 (a) at $t = 0$ are plotted in figure 2 (a) for $N_1^{(0)} = N_2^{(0)} = 1600$, $n_{11} = -0.000975$, $n_{22} = -0.000475$, $n_{12} = n_{21} = -0.000125$, $c = 1/16$, $\eta = 0.00625$, $x_{\text{max}} = 15$, with $\xi_i = \gamma_i = 0$. In the case of attractive interaction, for BEC with ^7Li the typical number of atoms in the

condensate is 10^3 and $|a_i|/a_{\text{ho}} \sim 10^{-3}$ to 10^{-4} [3]. In this example, $|a_1|/a_{\text{ho}} \simeq 3.5 \times 10^{-4}$, and $|a_2|/a_{\text{ho}} \simeq 1.7 \times 10^{-4}$. Hence the present numbers are well within the experimental scenario of BEC for atoms with attractive interaction. From figure 2 (a) we find that the wave-function components are more centrally peaked compared to those in figure 1 (a). This corresponds to small rms radii and large central density denoting an approximation to collapse.

Although the collapse of the coupled condensates could be inferred from the wave functions of figure 2 (a), we also study the dynamics of the collapse from a time evolution of GP equations (6) and (7) for $t > 0$ in the presence of an absorption and three-body recombination, e.g., for $\gamma_i \neq 0$ and $\xi_i \neq 0$ [4]. We consider the solution of (6) and (7) as shown in figure 2 (a) at $t = 0$ and allow this solution to evolve in time with $\gamma_i \neq 0$ and $\xi_i \neq 0$ by iteration. The general nature of time evolution is independent of the actual values of γ_i and ξ_i employed, provided that a very small value for ξ_i ($\sim 10^{-6}$) and a relatively larger one for γ_i (~ 0.1) are chosen [4]. In actual calculation we took $\gamma_1 = 0.15, \gamma_2 = 0.08$ and $\xi_1 = 0.00000031, \xi_2 = 0.00000094$. The quantities N_{at} and N_{mol} show the succession of collapses in a transparent fashion and we plot them in figure 2 (b) vs. time t . These quantities increase with time due to the absorptive interaction (γ_i). Once they are larger than the respective critical numbers for collapse, the three-body recombination takes over and these quantities fall below the critical level and the system collapses. Then the absorptive term is operative and these quantities increase again resulting in a sequence of collapses. In figure 2 (c) we plot the rms radii x_{rms} of the atomic and molecular condensates of the model of figure 2 (b) undergoing collapse for $t > 0$. The rms radii of the components execute oscillation similar to those in the uncoupled case [4].

The collapse was studied under various situations. After numerical investigation we find if only the diagonal parameter n_{11} or n_{22} is attractive and all other n_{ij} 's are repulsive then it is possible to have a collapse of component 1 or 2. If in addition to one diagonal parameter n_{ii} , the nondiagonal parameters n_{12} and n_{21} are attractive, it is possible to have a collapse in both components. If both diagonal parameters n_{11} and n_{22} are attractive, one can have

collapse of both the components.

Instead of reporting all our studies, in the following we consider the special case of collapse in the case where the diagonal elements n_{11} and n_{22} are positive and the nondiagonal terms n_{12} and n_{21} are negative. This corresponds to repulsion between two atoms and between two molecules and an attraction between a molecule and an atom. This case is of relevance, as experimentally, the rubidium system has been measured to have strongly attractive atom-molecule interaction and repulsive atom-atom and molecule-molecule interactions [2,7,10]. Also, it is now possible to vary continuously the nature (attractive or repulsive) and strength of interaction between bosons [24] which can create a specific situation of atom-molecule, atom-atom and molecule-molecule interactions. We find that when n_{12} and n_{21} are sufficiently attractive, they can surplus the repulsive terms n_{11} and n_{22} in (6) and (7) and the overall effective contribution of the quadratic nonlinear terms in these equations becomes attractive and one can have a collapse of both the components. To illustrate this interesting effect we study numerically the collapse for a negative n_{12} and n_{21} and positive n_{11} and n_{22} . In figure 3 (a) we plot the wave-function components obtained by solving (6) and (7) with $N_1^{(0)} = N_2^{(0)} = 1600$, $n_{11} = 0.0005$, $n_{22} = 0.000625$, $n_{12} = n_{21} = -0.00144$, $c = 1/16$, $\eta = 0.00625$, $x_{\max} = 10$ and $\gamma_i = \xi_i = 0$. Both wave-function components are peaked near $x = 0$ and have small rms radii. The system collapses with a small increase in $|n_{12}|$ and/or $|n_{21}|$. The time evolution of N_{at} and N_{mol} , and the corresponding rms radii are found by solving (6) and (7) iteratively with nonzero ξ_i and γ_i starting with the wave function presented in figure 3 (a) at $t = 0$. For $t > 0$ we employ $\xi_1 = 0.00000031$, $\xi_2 = 0.00000075$ and $\gamma_1 = 0.12$, $\gamma_2 = 0.05$. The quantities N_{at} and N_{mol} show the dynamics of collapse directly and they are plotted in figure 3 (b). The successive growth and decay of these numbers with time exhibit the sequence of collapses. Finally, in figure 3 (c) we plot the evolution of the corresponding rms radii. The results for $0 < t < 100$ in figures 2 and 3 are calculated with 2000 iterations of the GP equation (6) and (7) using a time step 0.05.

V. CONCLUSION

To conclude, we studied the stability and collapse of a trapped hybrid BEC consisting of atoms and molecules [6,7,9,10] coupled by coherent Raman transition [16] or Feshbach resonance [8,17] using the time-dependent coupled GP equation. An additional nonlinear interaction between the atoms and molecules given by (1) is introduced in the formulation which allows spontaneous transformation of two atoms into a molecule and vice versa. The effect of an external source of boson on the condensate is simulated by the absorptive term γ_i and the effect of three-body recombination necessary for the study of collapse is included via the quartic nonlinear term ξ_i in the GP equation.

We find interesting oscillation of the number of atoms and molecules due to the transformation of atoms to molecules and vice versa even when all interactions are repulsive. The component i of the condensate could experience collapse when the interaction among bosons of type i is attractive. Both components could experience collapse when at least the interaction between a molecule and an atom is attractive. They could also experience collapse when two of the bosonic interactions are attractive. The time evolution of collapse is studied via the time-dependent GP equation with absorption and three-body recombination. The number of particles of the component(s) of BEC experiencing collapse alternately grows and decays with time. The rms radii of the collapsing condensate also exhibit interesting oscillation as in the uncoupled case [4]. The oscillation of the rms radii denotes a radial oscillation of the condensates which can be observed experimentally. Recently, we studied the possibility of collapse in a coupled atomic BEC without any interaction allowing for transformation of one type of atom to another [25]. With the possibility of observation of coupled BEC, the findings of this study and of ref. [25] could be verified experimentally in the future.

We thank Dr. Kh. F. Abdullaev for a brief discussion. The work is supported in part by the Conselho Nacional de Desenvolvimento Científico e Tecnológico and Fundação de Amparo à Pesquisa do Estado de São Paulo of Brazil.

REFERENCES

- [1] Ensher J R, Jin D S, Matthews M R, Wieman C E and Cornell E A 1996 *Phys. Rev. Lett.* **77** 4984
- Davis K B, Mewes M O, Andrews M R, van Druten N J, Durfee D S, Kurn D M and Ketterle W 1995 *Phys. Rev. Lett.* **75** 3969
- Fried D G, Killian T C, Willmann L, Landhuis D, Moss S C, Kleppner D and Greytak T J 1998 *Phys. Rev. Lett.* **81** 3811
- [2] Wynar R, Freeland R S, Han D J, Ryu C and Heinzen D J 2000 *Science* **287** 1016
- [3] Bradley C C, Sackett C A, Tolett J J and Hulet R G 1995 *Phys. Rev. Lett.* **75** 1687
- Sackett C A, Stoof H T C and Hulet R G 1998 *Phys. Rev. Lett.* **80** 2031
- [4] Kagan Yu, Muryshv A E and Shlyapnikov G V 1998 *Phys. Rev. Lett.* **81** 933
- Gammal A, Frederico T , Tomio L and Chomaz Ph 2000 *Phys. Rev. A* **61** 051602
- [5] Ruprecht P A, Holland M J, Burnett K and Edwards M 1995 *Phys. Rev. A* **51** 4704
- Baym G and C.J. Pethick C J 1996 *Phys. Rev. Lett.* **76** 6
- Dalfovo F and Stringari S 1996 *Phys. Rev. A* **53** 2477
- Kim Y E and Zubarev A L 1998 *Phys. Lett. A* **246** 389
- Ueda M and Leggett A J 1998 *Phys. Rev. Lett.* **80** 1576 (1998)
- Wadati M and Tsurumi T 1998 *Phys. Lett. A* **247** 287
- Ueda M and Huang K 1999 *Phys. Rev. A* **61** 043601
- Saito H and Ueda M 2000 *Phys. Rev. Lett.* **86** 1406
- Saito H and Ueda M 2001 *Phys. Rev. A* **63** 043601
- Adhikari S K 2001 *Phys. Lett. A* **281** 265
- Abdullaev F Kh, Gammal A, Tomio L and Frederico T 2001 *Phys. Rev. A* **63** 043604

- Gammal A, Frederico T, Tomio L and Abdullaev F Kh 2000 *Phys. Lett. A* **267** 305
- Fetter A L 1995 cond-mat/9510037 unpublished
- [6] Drummond P D, Kheruntsyan K V and He H 1998 *Phys. Rev. Lett.* **81** 3055
- [7] Heinzen D J, Wynar R, Drummond P D and Kheruntsyan K V 2000 *Phys. Rev. Lett.* **84** 5029
- [8] Timmermans E, Tommasini P, Cote R, Hussein M and Kerman A 1999 *Phys. Rev. Lett.* **83** 2691
- Timmermans E, Tommasini P, Hussein M and Kerman A 1999 *Phys. Rep.* **315** 199
- [9] Góral K, Gaida M and Rzażewski K 2001 *Phys. Rev. Lett.* **86** 1397
- [10] Cusack B J, Alexander T J, Ostrovskaya E A and Kivshar Y S, 2001 *Phys. Rev. A* **64** 0236xx
- [11] Dalfovo F, Giorgini S, Pitaevskii L P and Stringari S 1999 *Rev. Mod. Phys.* **71** 463, see specially page 474.
- [12] Giorgini S, Pitaevskii L P and Stringari S 1996 *Phys. Rev. A* **54** R4633
- Edwards M, Ruprecht P A, Burnett K, Dodd R J and Clark C W 1996 *Phys. Rev. Lett.* **77** 1671
- Adhikari S K and Gammal A 2000 *Physica A* **286** 299
- [13] Adhikari S K 2000 *Phys. Lett. A* **265** 91
- Adhikari S K 2000 *Phys. Rev. E* **62** 2937
- Adhikari S K 2001 *Phys. Rev. E* **63** 056704
- [14] Dodd R J, Edwards M, Williams C J, Clark C W, Holland M J, Ruprecht P A and Burnett K 1996 *Phys. Rev. A* **54** 661
- Houbiers M and Stoof H T C 1996 *Phys. Rev. A* **54** 5055

- Eleftheriou A and Huang K 2000 *Phys. Rev. A* **61** 043601
- Adhikari S K 2000 *Physica A* **284** 97
- [15] Gross E P 1961 *Nuovo Cimento* **20** 454
- Pitaevskii L P 1961 *Zh. Eksp. Teor. Fiz.* **40** 646 [1961 *Sov. Phys. JETP* **13** 451]
- [16] Fioretti A, Comparat D, Crubellier A, Dulieu O, Masnou-Seeuws F and Pillet P 1998 *Phys. Rev. Lett.* **80** 4402
- Julienne P S, Burnett K, Band Y B and Stwalley W C 1998 *Phys. Rev. A* **58** R797
- [17] Courteille Ph, Freeland R S, Heinzen D J, van Abeelen F A and Verhaar B J 1998 *Phys. Rev. Lett.* **81** 69
- Inouye S, Andrews M R, Stenger J, Miesner H J, Stamper-Kurn D M and Ketterle W 1998 *Nature (London)* **392** 151
- [18] Matthews M R, Anderson B P, Haljan P C, Hall D S, Wieman C E and Cornell E A 1999 *Phys. Rev. Lett.* **83** 2498
- Matthews M R, Hall D S, Jin D S, Ensher J R, Wieman C E, Cornell E A, Dalfovo F, Minniti C and Stringari S 1998 *Phys. Rev. Lett.* **81** 243
- [19] Goldstein E V and Meystre P 1997 *Phys. Rev. A* **55** 2935
- Pu H and Bigelow N P 1998 *Phys. Rev. Lett.* **80** 1134
- [20] Tolmachev V V 2000 *Phys. Lett. A* **266** 400
- Friedberg R and Lee T D 1989 *Phys. Rev. B* **40** 6745
- [21] Tsai C C, Freeland R S, Vogels J M, Boesten H M J M, Verhaar B J and Heinzen D J 1997 *Phys. Rev. Lett.* **79** 1245
- [22] Sinatra A, Fedichev P O, Castin Y, Dalibard J and Shlyapnikov G V 1999 *Phys. Rev. Lett.* **82** 251

- Esry B D, Greene C H, Burke Jr J P and Bohn J L 1997 *Phys. Rev. Lett.* **78** 3594
- Ho T-L and Shenoy V B 1996 *Phys. Rev. Lett.* **77** 3276
- Dalfovo F and Modugno M 2000 *Phys. Rev. A* **61** 023605
- Esry B D 1998 *Phys. Rev. A* **58** R3399
- [23] Koonin S E 1986 *Computational Physics* (Melno Park: Benjamin/Cummings Pub. Co. Inc.) pp 161-167.
- [24] J. L. Roberts, N. R. Claussen, S. L. Cornish, E. A. Donley, E. A. Cornell, C. E. Wieman, *Phys. Rev. Lett.* **86**, 4211 (2001).
- [25] Adhikari S K 2001 *Phys. Rev. A* **63** 043611

Figure Caption:

1. (a) Atomic and molecular wave function components $\phi_1(x)$ and $\phi_2(x)$ vs. x at $t = 0$, (b) the corresponding numbers N_{at} and N_{mol} vs. time t (> 0), and (c) the rms radii vs. time t (> 0) obtained for $N_1^{(0)} = N_2^{(0)} = 10000$, $n_{11} = n_{22} = 0.001$, $n_{12} = n_{21} = 0.0005$, $c = 1/16$, for $\xi_i = \gamma_i = 0$. The curves are labelled by the corresponding values of η .

2. The same as in figure 1 with $N_1^{(0)} = N_2^{(0)} = 1600$, $n_{11} = -0.000975$, $n_{22} = -0.000475$, $n_{12} = n_{21} = -0.000125$, $c = 1/16$, and $\eta = 0.00625$. In (a) $\xi_i = \gamma_i = 0$, and in (b) and (c) $\xi_1 = 0.00000031$, $\gamma_1 = 0.15$, $\xi_2 = 0.00000094$, $\gamma_2 = 0.08$.

3. The same as in figure 1 with $N_1^{(0)} = N_2^{(0)} = 1600$, $n_{11} = 0.0005$, $n_{22} = 0.000625$, $n_{12} = n_{21} = -0.00144$, $c = 1/16$, and $\eta = 0.00625$. In (a) $\xi_i = \gamma_i = 0$, and in (b) and (c) $\xi_1 = 0.00000031$, $\gamma_1 = 0.12$, $\xi_2 = 0.00000075$, $\gamma_2 = 0.05$.

Figure 1 (a)




Figure 1 (b)




Figure 1 (c)




Figure 2 (a)




Figure 2 (b)




Figure 2 (c)




Figure 3 (a)




Figure 3 (b)




Figure 3 (c)

